

Probabilistic chain teleportation of a qutrit-state

Meiyu Wang, Fengli Yan*

*College of Physics Science and Information Engineering,
Hebei Normal University, Shijiazhuang 050016, China
Hebei Advanced Thin Films Laboratory,
Shijiazhuang 050016, China
(Dated: March 25, 2010)*

We investigate chain teleportation of a qutrit-state via the non-maximally two-qutrit entangled channels. For the case of four parties, the efficiencies of two chain teleportation protocols, the separate chain teleportation protocol (SCTP) and the global chain teleportation protocol (GCTP), are calculated. In SCTP the errors are corrected between every step while in GCTP the errors are corrected only at the end. Furthermore, we present a piecewise global chain teleportation protocol (PGCTP) for keeping away from the inconvenience of error-correction of GCTP. We show that PGCTP is more efficient than SCTP.

PACS numbers: 03.67.Hk

Quantum teleportation plays an important role in the field of quantum computation and quantum communication. It allows a sender (called Alice) to transmit an unknown quantum state to a receiver (called Bob) by using an entangled state as a quantum resource and by sending classical information via ordinary channel. The original protocol of Bennett et al. [1] involves teleportation of an arbitrary state of a qubit via an Einstein-Podolsky-Rosen (EPR) pair and by transmitting two bits of classical information from Alice to Bob. Here Alice knows neither the state to be teleported nor the location of the intended receiver, Bob. They also presented a protocol for teleporting an unknown state of a qudit via a maximally entangled state in $d \times d$ dimensional Hilbert space and by sending $2 \log_2 d$ bits of classical information. Since then, quantum teleportation has got great development [2–16] and has been experimentally demonstrated by several groups [17–20].

However in the real world situations, it is most of the time not possible to have a maximally entangled state at one's disposal. Because of the interaction with the environment, the state of any system would become a mixed state after a certain period. This problem of decoherence can be mitigated but cannot be completely overcome easily. Also, it may happen that the source does not produce perfect EPR pairs rather non-maximally entangled pairs which is shared between Alice and Bob. If we have non-maximally entangled state as a shared resource and we want to do quantum teleportation, we have to pay some price. That is, we have to compromise either in fidelity or in the success probability. If we are ready to pay the price for the success probability it is possible to have unit fidelity teleportation. And this scheme is called probabilistic quantum teleportation [12, 13].

When one considers teleportation in a quantum chain network, the teleportation strategy is miscellaneous.

Suppose that the quantum chain network consists of $N+1$ parties (called Alice1, Alice2, Alice3, ..., Alice $N+1$ respectively in sequence) who share the entangled sources between the neighboring parties. The simplest and most direct strategy is the separate chain teleportation protocol (SCTP) by performing N separate teleportations, i.e., Alice1 teleports the quantum state to Alice2 via the first teleportation. Then Alice2 teleports it to Alice3 via the second teleportation and so on. If the entangled sources between every neighboring parties are the same, the total success probability equals p^N when a quantum state teleportation from starting point Alice1 to end point Alice $N+1$, where p is the success probability of one step. The total success probability of SCTP is very small for $p < 1$ when N is large. A more efficient strategy is the global chain teleportation protocol (GCTP) [14]. In an interesting work, Modlowska and Grudka [14] showed that if the qubit is teleported several times via some non-maximally entangled states, the “errors” introduced in the previous teleportations can be corrected by the “errors” introduced in the following teleportations. This effect is called error self-correction. The total success probability would be enhanced immensely if we correct all errors in the final receiver. Their strategy was developed in the framework of the scheme proposed in Ref. [4] for linear optical teleportation. In the recent papers [15, 16], it has been shown that this feature of the multiple teleportation of Ref. [14] is not restricted to the teleportation scheme stated in Ref. [4]. In Ref. [16], we present global chain teleportation protocol based on the general teleportation language of the original proposal shown in Ref. [1].

In this paper, we would investigate the probabilistic chain teleportation of qutrit-state via non-maximally two-qutrit entangled channels. For the case of four parties, the efficiencies of two chain teleportation protocols, the SCTP and the GCTP, are calculated. With the increase of the parties number N , the collapsing states will become more and more complex if we apply the GCTP. For keeping away from this inconvenience, we present the

*Electronic address: flyan@mail.hebtu.edu.cn

piecewise global chain teleportation protocol (PGCTP). We show that PGCTP is more efficient than SCTP.

It is easy to see that the error self-correction only appears in the case of $3N + 1$ parties for chain teleportation of qutrit state. Therefore, to illustrate our protocol clearly, let us first begin with the chain teleportation in the case of four parties.

Alice wants to teleport an unknown quantum state

$$|\psi\rangle = \alpha|0\rangle + \beta|1\rangle + \gamma|2\rangle \quad (1)$$

to Bob, where α, β and $\gamma \in \mathbb{C}$ and $|\alpha|^2 + |\beta|^2 + |\gamma|^2 = 1$. There is no direct entangled resource between Alice and Bob. Fortunately, Alice and Bob can build the relationship by two intermediaries Charlie and Dave. Alice and Charlie share a non-maximally two-qutrit entangled resource

$$|\Psi\rangle = a_0|00\rangle + a_1|11\rangle + a_2|22\rangle, \quad (2)$$

while Charlie and Dave, Dave and Bob share the identical resource, where a_0, a_1 and a_2 are real numbers and satisfy $a_0^2 + a_1^2 + a_2^2 = 1$. Without loss of generality, we suppose $a_0 \leq a_1 \leq a_2$.

Firstly, we consider the SCTP completed by performing three separate standard teleportations, i.e., Alice teleports the quantum state $|\psi\rangle$ to Charlie via the first teleportation. Then Charlie teleports it to Dave via the second teleportation. Finally, Dave teleports it to receiver Bob via the third teleportation. According to the standard probabilistic teleportation protocol, in the first separate teleportation, Alice performs generalized Bell-basis measurement (GBM) in the basis $\{|\Phi_{mn}\rangle, m, n = 0, 1, 2\}$ on the teleported qutrit and the entangled qutrit in her side.

$$|\Phi_{mn}\rangle = \frac{1}{\sqrt{3}} \sum_{j=0}^2 e^{2ijn\pi/3} |j\rangle |(j \oplus m)\rangle, \quad (3)$$

where here and hereafter $j \oplus m$ means sum of j and m modulo 3. Charlie can apply the corresponding unitary transformation conditioned on the result of Alice's measurement. The corresponding unitary transformations read

$$\begin{aligned} U_{00} &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, U_{01} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & e^{-2\pi i/3} & 0 \\ 0 & 0 & e^{-4\pi i/3} \end{bmatrix}, \\ U_{02} &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & e^{-4\pi i/3} & 0 \\ 0 & 0 & e^{-2\pi i/3} \end{bmatrix}, \\ U_{10} &= \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, U_{11} = \begin{bmatrix} 0 & 0 & e^{-4\pi i/3} \\ 1 & 0 & 0 \\ 0 & e^{-2\pi i/3} & 0 \end{bmatrix}, \\ U_{12} &= \begin{bmatrix} 0 & 0 & e^{-2\pi i/3} \\ 1 & 0 & 0 \\ 0 & e^{-4\pi i/3} & 0 \end{bmatrix}, \end{aligned}$$

$$\begin{aligned} U_{20} &= \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}, U_{21} = \begin{bmatrix} 0 & e^{-2\pi i/3} & 0 \\ 0 & 0 & e^{-4\pi i/3} \\ 1 & 0 & 0 \end{bmatrix}, \\ U_{22} &= \begin{bmatrix} 0 & e^{-4\pi i/3} & 0 \\ 0 & 0 & e^{-2\pi i/3} \\ 1 & 0 & 0 \end{bmatrix}. \end{aligned} \quad (4)$$

Finally, the state Bob received becomes

$$|\psi_1\rangle = \frac{1}{\sqrt{p_1}} (\alpha a_0|0\rangle + \beta a_1|1\rangle + \gamma a_2|2\rangle) \quad (5)$$

with the probability $p_1 = |a_0\alpha|^2 + |a_1\beta|^2 + |a_2\gamma|^2$ or

$$|\psi_2\rangle = \frac{1}{\sqrt{p_2}} (\alpha a_1|0\rangle + \beta a_2|1\rangle + \gamma a_0|2\rangle) \quad (6)$$

with the probability $p_2 = |a_1\alpha|^2 + |a_2\beta|^2 + |a_0\gamma|^2$ or

$$|\psi_3\rangle = \frac{1}{\sqrt{p_3}} (\alpha a_2|0\rangle + \beta a_0|1\rangle + \gamma a_1|2\rangle) \quad (7)$$

with the probability $p_3 = |a_2\alpha|^2 + |a_0\beta|^2 + |a_1\gamma|^2$.

These states are in accordance with the original state $|\psi\rangle$ only if the quantum channel is a maximally entangled state, i.e. $a_0 = a_1 = a_2$. For the case of non-maximally entangled channel, these states can be returned to the original state with certain probability by performing the generalized measurement given by Kraus operators:

$$E_{S1} = |0\rangle\langle 0| + \frac{a_0}{a_1}|1\rangle\langle 1| + \frac{a_0}{a_2}|2\rangle\langle 2|, \quad (8a)$$

$$E_{F1} = \sqrt{1 - \frac{a_0^2}{a_1^2}}|1\rangle\langle 1| + \sqrt{1 - \frac{a_0^2}{a_2^2}}|2\rangle\langle 2| \quad (8b)$$

for $|\psi_1\rangle$ and

$$E_{S2} = \frac{a_0}{a_1}|0\rangle\langle 0| + \frac{a_0}{a_2}|1\rangle\langle 1| + |2\rangle\langle 2|, \quad (9a)$$

$$E_{F2} = \sqrt{1 - \frac{a_0^2}{a_1^2}}|0\rangle\langle 0| + \sqrt{1 - \frac{a_0^2}{a_2^2}}|1\rangle\langle 1| \quad (9b)$$

for $|\psi_2\rangle$ and

$$E_{S3} = \frac{a_0}{a_2}|0\rangle\langle 0| + |1\rangle\langle 1| + \frac{a_0}{a_1}|2\rangle\langle 2|, \quad (10a)$$

$$E_{F3} = \sqrt{1 - \frac{a_0^2}{a_2^2}}|0\rangle\langle 0| + \sqrt{1 - \frac{a_0^2}{a_1^2}}|2\rangle\langle 2| \quad (10b)$$

for $|\psi_3\rangle$.

When E_S is obtained, the qutrit ends in its original state $|\psi\rangle = \alpha|0\rangle + \beta|1\rangle + \gamma|2\rangle$. The success probability in the first teleportation is

$$p = \sum_{i=1}^3 p_i \langle \psi_i | E_{Si}^\dagger E_{Si} | \psi_i \rangle = 3a_0^2. \quad (11)$$

Next, Charlie teleports the recovered quantum state to Dave, and Dave teleports the recovered quantum state

to Bob by the similar process. Combining these three teleportations, the total probability that Bob receives the quantum state $|\psi\rangle$ is

$$P_S(4) = p^3 = 27a_0^6. \quad (12)$$

However, the above teleportation protocol is not the optimal strategy. In fact, the intermediaries Charlie and Dave does not need to recover the quantum state to be teleported. They only make the conditioned unitary transformation according to the previous party's measurement outcome and teleports the "errors" state to the next party directly. Lastly, Bob corrects all "errors" of the quantum state in the teleportation process. This is so called GCTP.

Let us, thus, assume that Charlie and Dave does not recover the quantum state to be teleported in the teleportation, they only make a unitary transformation according to the previous party's measurement outcome, then they also perform GBM on their two qutrits and broadcast the measurement outcome to the next party. After making the corresponding transformation conditioned on Dave's measurement outcome, Bob's qutrit will collapse into one of the following states

$$|\phi_1\rangle = a_0^3\alpha|0\rangle + a_1^3\beta|1\rangle + a_2^3\gamma|2\rangle, \quad (13a)$$

$$|\phi_2\rangle = a_1^3\alpha|0\rangle + a_2^3\beta|1\rangle + a_0^3\gamma|2\rangle, \quad (13b)$$

$$|\phi_3\rangle = a_2^3\alpha|0\rangle + a_0^3\beta|1\rangle + a_1^3\gamma|2\rangle, \quad (13c)$$

$$|\phi_4\rangle = a_0^2a_1\alpha|0\rangle + a_1^2a_2\beta|1\rangle + a_2^2a_0\gamma|2\rangle, \quad (13d)$$

$$|\phi_5\rangle = a_1^2a_2\alpha|0\rangle + a_2^2a_0\beta|1\rangle + a_0^2a_1\gamma|2\rangle, \quad (13e)$$

$$|\phi_6\rangle = a_2^2a_0\alpha|0\rangle + a_0^2a_1\beta|1\rangle + a_1^2a_2\gamma|2\rangle, \quad (13f)$$

$$|\phi_7\rangle = a_0^2a_2\alpha|0\rangle + a_1^2a_0\beta|1\rangle + a_2^2a_1\gamma|2\rangle, \quad (13g)$$

$$|\phi_8\rangle = a_1^2a_0\alpha|0\rangle + a_2^2a_1\beta|1\rangle + a_0^2a_2\gamma|2\rangle, \quad (13h)$$

$$|\phi_9\rangle = a_2^2a_1\alpha|0\rangle + a_0^2a_2\beta|1\rangle + a_1^2a_0\gamma|2\rangle, \quad (13i)$$

$$|\phi_{10}\rangle = \alpha|0\rangle + \beta|1\rangle + \gamma|2\rangle \quad (13j)$$

with the probabilities

$$\begin{aligned} p'_1 &= a_0^6|\alpha|^2 + a_1^6|\beta|^2 + a_2^6|\gamma|^2, \\ p'_2 &= a_1^6|\alpha|^2 + a_2^6|\beta|^2 + a_0^6|\gamma|^2, \\ p'_3 &= a_2^6|\alpha|^2 + a_0^6|\beta|^2 + a_1^6|\gamma|^2, \\ p'_4 &= 3(a_0^4a_1^2|\alpha|^2 + a_1^4a_2^2|\beta|^2 + a_2^4a_0^2|\gamma|^2), \\ p'_5 &= 3(a_1^4a_2^2|\alpha|^2 + a_2^4a_0^2|\beta|^2 + a_0^4a_1^2|\gamma|^2), \\ p'_6 &= 3(a_2^4a_0^2|\alpha|^2 + a_0^4a_1^2|\beta|^2 + a_1^4a_2^2|\gamma|^2), \\ p'_7 &= 3(a_0^4a_2^2|\alpha|^2 + a_1^4a_0^2|\beta|^2 + a_2^4a_1^2|\gamma|^2), \\ p'_8 &= 3(a_1^4a_0^2|\alpha|^2 + a_2^4a_1^2|\beta|^2 + a_0^4a_2^2|\gamma|^2), \\ p'_9 &= 3(a_2^4a_1^2|\alpha|^2 + a_0^4a_2^2|\beta|^2 + a_1^4a_0^2|\gamma|^2), \\ p'_{10} &= 6a_0^2a_1^2a_2^2. \end{aligned}$$

respectively. When the state is in $|\phi_{10}\rangle$, we do not have to perform the error correction for the errors self-correction in the teleportation. For $|\phi_i\rangle$ ($i = 1, 2, \dots, 6$), one can

recover the original state by performing generalized measurement given by Kraus operators:

$$E'_{S1} = |0\rangle\langle 0| + \frac{a_0^3}{a_1^3}|1\rangle\langle 1| + \frac{a_0^3}{a_2^3}|2\rangle\langle 2|, \quad (14a)$$

$$E'_{F1} = \sqrt{1 - \frac{a_0^6}{a_1^6}}|1\rangle\langle 1| + \sqrt{1 - \frac{a_0^6}{a_2^6}}|2\rangle\langle 2| \quad (14b)$$

for $|\phi_1\rangle$ and

$$E'_{S2} = \frac{a_0^3}{a_1^3}|0\rangle\langle 0| + \frac{a_0^3}{a_2^3}|1\rangle\langle 1| + |2\rangle\langle 2|, \quad (15a)$$

$$E'_{F2} = \sqrt{1 - \frac{a_0^6}{a_1^6}}|0\rangle\langle 0| + \sqrt{1 - \frac{a_0^6}{a_2^6}}|2\rangle\langle 2| \quad (15b)$$

for $|\phi_2\rangle$ and

$$E'_{S3} = \frac{a_0^3}{a_2^3}|0\rangle\langle 0| + |1\rangle\langle 1| + \frac{a_0^3}{a_1^3}|2\rangle\langle 2|, \quad (16a)$$

$$E'_{F3} = \sqrt{1 - \frac{a_0^6}{a_2^6}}|0\rangle\langle 0| + \sqrt{1 - \frac{a_0^6}{a_1^6}}|2\rangle\langle 2| \quad (16b)$$

for $|\phi_3\rangle$ and

$$E'_{S4} = |0\rangle\langle 0| + \frac{a_0^2}{a_1a_2}|1\rangle\langle 1| + \frac{a_0a_1}{a_2^2}|2\rangle\langle 2|, \quad (17a)$$

$$E'_{F4} = \sqrt{1 - \frac{a_0^4}{a_1^2a_2^2}}|1\rangle\langle 1| + \sqrt{1 - \frac{a_0^2a_1^2}{a_2^4}}|2\rangle\langle 2| \quad (17b)$$

for $|\phi_4\rangle$ and

$$E'_{S5} = \frac{a_0^2}{a_1a_2}|0\rangle\langle 0| + \frac{a_0a_1}{a_2^2}|1\rangle\langle 1| + |2\rangle\langle 2|, \quad (18a)$$

$$E'_{F5} = \sqrt{1 - \frac{a_0^4}{a_1^2a_2^2}}|0\rangle\langle 0| + \sqrt{1 - \frac{a_0^2a_1^2}{a_2^4}}|1\rangle\langle 1| \quad (18b)$$

for $|\phi_5\rangle$ and

$$E'_{S6} = \frac{a_0a_1}{a_2^2}|0\rangle\langle 0| + |1\rangle\langle 1| + \frac{a_0^2}{a_1a_2}|2\rangle\langle 2|, \quad (19a)$$

$$E'_{F6} = \sqrt{1 - \frac{a_0^2a_1^2}{a_2^4}}|0\rangle\langle 0| + \sqrt{1 - \frac{a_0^4}{a_1^2a_2^2}}|2\rangle\langle 2| \quad (19b)$$

for $|\phi_6\rangle$ respectively.

For $|\phi_7\rangle, |\phi_8\rangle$ and $|\phi_9\rangle$, the recover operator are somewhat complicate because $a_0^2a_2 \leq a_1^2a_0$ for some entangled channels but $a_0^2a_2 \geq a_1^2a_0$ for other entangled channels. The recover operators take different form for the above two cases.

For the case of $a_0^2a_2 \leq a_1^2a_0$, the recover Kraus operators take the form:

$$E'_{S7} = |0\rangle\langle 0| + \frac{a_0a_2}{a_1^2}|1\rangle\langle 1| + \frac{a_0^2}{a_1a_2}|2\rangle\langle 2|, \quad (20a)$$

$$E'_{F7} = \sqrt{1 - \frac{a_0^2a_2^2}{a_1^4}}|1\rangle\langle 1| + \sqrt{1 - \frac{a_0^4}{a_1^2a_2^2}}|2\rangle\langle 2| \quad (20b)$$

for $|\phi_7\rangle$ and

$$E'_{S8} = \frac{a_0 a_2}{a_1^2} |0\rangle\langle 0| + \frac{a_0^2}{a_1 a_2} |1\rangle\langle 1| + |2\rangle\langle 2|, \quad (21a)$$

$$E'_{F8} = \sqrt{1 - \frac{a_0^2 a_2^2}{a_1^4}} |0\rangle\langle 0| + \sqrt{1 - \frac{a_0^4}{a_1^2 a_2^2}} |1\rangle\langle 1| \quad (21b)$$

for $|\phi_8\rangle$ and

$$E'_{S9} = \frac{a_0^2}{a_1 a_2} |0\rangle\langle 0| + |1\rangle\langle 1| + \frac{a_0 a_2}{a_1^2} |2\rangle\langle 2|, \quad (22a)$$

$$E'_{F9} = \sqrt{1 - \frac{a_0^4}{a_1^2 a_2^2}} |0\rangle\langle 0| + \sqrt{1 - \frac{a_0^2 a_2^2}{a_1^4}} |2\rangle\langle 2| \quad (22b)$$

for $|\phi_9\rangle$ respectively.

On the other hand, if $a_0^2 a_2 \geq a_1^2 a_0$, the recover Kraus operators take the form:

$$E''_{S7} = \frac{a_1^2}{a_0 a_2} |0\rangle\langle 0| + |1\rangle\langle 1| + \frac{a_0 a_1}{a_2^2} |2\rangle\langle 2|, \quad (23a)$$

$$E''_{F7} = \sqrt{1 - \frac{a_1^4}{a_0^2 a_2^2}} |0\rangle\langle 0| + \sqrt{1 - \frac{a_0^2 a_1^2}{a_2^4}} |2\rangle\langle 2| \quad (23b)$$

for $|\phi_7\rangle$

$$E''_{S8} = |0\rangle\langle 0| + \frac{a_0 a_1}{a_2^2} |1\rangle\langle 1| + \frac{a_1^2}{a_0 a_2} |2\rangle\langle 2|, \quad (24a)$$

$$E''_{F8} = \sqrt{1 - \frac{a_0^2 a_1^2}{a_2^4}} |1\rangle\langle 1| + \sqrt{1 - \frac{a_1^4}{a_0^2 a_2^2}} |2\rangle\langle 2| \quad (24b)$$

for $|\phi_8\rangle$ and

$$E''_{S9} = \frac{a_0 a_1}{a_2^2} |0\rangle\langle 0| + \frac{a_1^2}{a_0 a_2} |1\rangle\langle 1| + |2\rangle\langle 2|, \quad (25a)$$

$$E''_{F9} = \sqrt{1 - \frac{a_0^2 a_1^2}{a_2^4}} |0\rangle\langle 0| + \sqrt{1 - \frac{a_1^4}{a_0^2 a_2^2}} |1\rangle\langle 1| \quad (25b)$$

for $|\phi_9\rangle$ respectively.

The probability of successfully recovering the original state is

$$P_G(4) = 3a_0^6 + 9a_0^4 a_1^2 + 9 \min\{a_0^4 a_2^2, a_1^4 a_0^2\} + 6a_0^2 a_1^2 a_2^2. \quad (26)$$

We can easily see $P_G(4) \geq P_S(4)$ because of $a_2 \geq a_1 \geq a_0$. It is obvious that for the maximally entangled channel, the two protocols are equivalent, but for the non-maximally entangled channel, GCTP is more efficient than SCTP. The total success probability of SCTP is only determined by the smallest coefficient of the entangled channel, on the other hand, the total success probability of GCTP lies on all coefficients of the entangled channel. For fixed a_0 , the total success probability of GCTP takes the minimum $P_G^{\min}(4) = 6a_0^4 + 9a_0^6$

for the case of $a_1 = a_0$ and the maximum $P_G^{\max}(4) = \frac{3}{2}a_0^2 + 6a_0^4 - \frac{9}{2}a_0^6$ for the case of $a_1 = a_2$.

When one wants to apply global chain teleportation of a qutrit for the case of arbitrary parties, a hindrance occurs. With the increase of the party number, the possible collapsing states of receiver's qutrit will become numerous and disorderly. So the error-correction operation of GCTP is very inconvenient. But we can select a teleportation strategy eclectically between GCTP and SCTP. Instead of being corrected by the final receiver in GCTP and corrected by every parties in SCTP, the errors can be corrected every n parties (n is small). We call it piecewise globe chain teleportation protocol (PGCTP). Here we select $n = 3$ since error self-correction only appears after making 3 times generalized Bell-basis measurements.

Suppose that Alice1 wants to teleport a quantum state $|\psi\rangle = \alpha|0\rangle + \beta|1\rangle + \gamma|2\rangle$ to Alice $3N + 1$. There is no direct entangled resource between them, but they can link through $3N - 1$ intermediaries called Alice2, Alice3, $\dots$, Alice $3N$, respectively. Two neighboring parties share the partially entangled state described by Eq.(2).

Firstly, Alice1, Alice2, Alice3 and Alice4 apply the GCTP according to the steps as above described. After correct all errors, Alice4 obtain the quantum state $|\psi\rangle$ with the probability $P_G(4)$. Next, Alice4, Alice5, Alice6 and Alice7 apply the GCTP again. Thus Alice7 obtains the state $|\psi\rangle$ with the probability $P_G(4)^2$, and so on. Finally, Alice $3N + 1$ obtain the quantum state $|\psi\rangle$ with the probability

$$P_{PG}(3N + 1) = P_G(4)^N \quad (27)$$

after correct the errors.




FIG. 1: The total success probability P_{PG} and P_S versus a_0 for different N (Solid line: P_S , shadow region: P_{PG} , where upper contour line corresponding to the case of $a_1 = a_2$, and lower contour line corresponding to the case of $a_0 = a_1$). From (a) to(d), N takes 1, 2, 5, 10 accordingly.

In Fig.1, we plot P_S and P_{PG} as the function of a_0

for different N . We can see that both the total success probabilities of two protocols declines with the decrease of the entanglement of channels. Moreover, the greater N is, the more sharper the success probabilities declines. It shows that the quantum channel with small entanglement will become unpractical with the increase of N . Fig.1 also indicates explicitly that the PGCTP is more efficient than SCTP. For example, for the case of $N = 5$, the total success probability of PGCTP $P_{PG}^{\max} \approx 15\%$ while the total success probability of SCTP P_S only attains 1.3% when the smallest coefficient of channels $a_0 = 0.50$.




FIG. 2: The ratio of P_{PG} to P_S versus a_0 for different N : upper contour line corresponding to the case of $a_1 = a_2$, and lower contour line corresponding to the case of $a_0 = a_1$. From (a) to (d), N takes 1, 2, 5, 10 accordingly.

The ratio of P_{PG} to P_S as a function of a_0 for different N is illustrated in Fig.2. Here we only take a_0 from 0.5 to $\frac{1}{\sqrt{3}}$ because the small entanglement channels are unpractical for large N . From Fig.2, we can see that the greater N is, the larger P_{PG}/P_S is. In other words, the efficiency of PGCTP is far higher than that of SCTP when the steps of teleportation increase.

In summary, we have studied chain teleportation of a qutrit-state via the non-maximally two-qutrit entangled channels. For the case of four parties, the efficiencies of two chain teleportation protocols, the separate chain teleportation protocol and the global chain teleportation protocol, are calculated. In SCTP the errors are corrected between every step while in GCTP the errors are corrected only at the end. With the increase of the parties number, possible collapsing states of receiver's qutrit will become numerous and disorderly. Therefore, the error-corrections of GCTP become very inconvenient. We present a piecewise global chain teleportation protocol for keeping away from this inconvenience. We show that PGCTP is more efficient than SCTP.

Acknowledgments

This work was supported by the National Natural Science Foundation of China under Grant No: 10971247, Hebei Natural Science Foundation of China under Grant No: F2009000311 and the Key Project of Science and Technology Research of Education Ministry of China under Grant No: 207011.

-
- [1] C.H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W.K. Wootters, Phys. Rev. Lett. **70**, 1895 (1993).
 - [2] S.L. Braunstein and P. van Loock, Rev. Mod. Phys. **77**, 427 (2005).
 - [3] X.B. Wang, T. Hiroshima, A. Tomita, and M. Hayashi, Phys. Rep. **448**, 1 (2007).
 - [4] E. Knill, R. Laflamme, and G.J. Milburn, Nature (London) **409**, 46 (2001).
 - [5] L. Vaidman, Phys. Rev. A **49**, 1473 (1994).
 - [6] S.L. Braunstein and H.J. Kimble, Phys. Rev. Lett. **80**, 869 (1998).
 - [7] G. Gordon and G. Rigolin, Phys. Rev. A **73**, 042309 (2006).
 - [8] A. Karlsson and M. Bourennane, Phys. Rev. A **58**, 4394 (1998).
 - [9] F.G. Deng, C.Y. Li, Y.S. Li, H.Y. Zhou, and Y. Wang, Phys. Rev. A **72**, 022338 (2005).
 - [10] Z.J. Zhang, Phys. Lett. A **352**, 55 (2006).
 - [11] T. Gao, F.L. Yan, and Y.C. Li, Europhys. Lett. **84**, 50001 (2008).
 - [12] P. Agrawal and A.K. Pati, Phys. Lett. A **305**, 12 (2002).
 - [13] A.K. Pati and P. Agrawal, J. Opt. B: Quantum Semiclassical Opt. **6**, S844 (2004).
 - [14] J. Modlowska and A. Grudka, Phys. Rev. Lett. **100**, 110503 (2008).
 - [15] G. Rigolin, J. Phys. B: At. Mol. Opt. Phys. **42**, 235504 (2009).
 - [16] M. Y. Wang and F. L. Yan, Euro. Phys. J. D **54**, 111 (2009).
 - [17] D. Bouwmeester, J.W. Pan, K. Mattle, M. Eibl, H. Weinfurter, and A. Zeilinger, Nature **390**, 575(1997).
 - [18] A. Furusawa, J.L. Sørensen, S.L. Braunstein, C.A. Fuchs, H.J. Kimble, and E.S. Polzik, Science **282**, 706 (1998).
 - [19] M.A. Nielsen, E. Knill, and R. Laflamme, Nature **396**, 52 (1998).
 - [20] D. Boschi, S. Branca, F. De Martini, L. Hardy, and S. Popescu, Phys. Rev. Lett. **80**, 1121 (1998).